



Pseudomonas Agar Base

(C.F.C./C.N. Agar)

LAB 108

Description

The base medium is a modification of King's medium A which uses magnesium and potassium salts to enhance production of the pigments pyocyanin (green) and fluorescein (detected by U.V./blue light). The medium is made selective for *Pseudomonas aeruginosa* by the addition of X107 C.N. supplement. Alternatively the medium can be made selective for *Pseudomonas* species generally by the addition of X108 C.F.C. supplement. This medium can be made selective for the isolation of *Burkholderia cepacia* by the addition of X140.

Typical Formula	g/litre
Acid Hydrolysed Casein	10.0
Gelatin Peptone	16.0
Potassium sulphate	10.0
Magnesium chloride	1.4
Agar No. 2	11.0

Method for reconstitution

Weigh 48.4 grams of powder and disperse in 1 litre of deionised water. Add 10ml of glycerol. Sterilise by autoclaving at 121°C for 15 minutes. Allow the medium to cool to 47°C then add the contents of 2 vials of either X107 C.N. supplement or X108 C.F.C. supplement. Mix well and pour into Petri dishes.

Appearance: Pale straw, opaque.

pH: 7.1 ± 0.2

Minimum Q.C. organisms: *P. aeruginosa* WDCM 00025
E. coli (inhibition) WDCM 00013

Storage of Prepared Medium: Plates – up to 7 days at 2-8°C in the dark.

Inoculation: Surface, spread 0.1 to 0.5ml of sample over entire surface.

Incubation: 25-30°C aerobically for 48 hours.

Interpretation: Count all colonies as *Pseudomonas species*. Colonies that exhibit the pyocyanin and fluorescein pigments count as *P. aeruginosa*.

Growth Characteristics				
organism	colony size (mm)	shape & surface	colour	fluorescence
<i>Ps. aeruginosa</i>	2.0-3.0	CV.Cr.D.	Green/Blue	yes
<i>P. fluorescens</i>	2.0-3.0	CV.Cr.D.	Yellow	yes
<i>P. fragi</i>	1.0-3.0	CV.Cr.D.	Grey	no

References

Burton, M.O., Campbell, J.J.R. and Eagles, B.A. (1948). The mineral requirement for pyocyanin production. Can. J. res. Sect. C. Bot. Sci. 26:15.

King, E.O., Ward, M.K. and Raney, D.E. (1954). Two simple media for the demonstration of pyocyanin and fluorescein. J. Lab. Clin. Med. 44: 301.

Goto, S. and Enomoto, S. (1970). Jap. J. Microbiol. 14: 65-72.

Mead, G.C. and Adams, B.W. (1977). Br. Poult. Sci. 18: 661-667.